



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore \rho = r \sqrt{\left(\frac{l \frac{n-n_r}{n}}{l \frac{1}{2}} \right)} = r \sqrt{\left(\frac{l \frac{n-n_r}{n}}{l \frac{1}{2}} \right)}. \quad (10)$$

This agrees with Herschel's formulæ which Prof. Chase employs.

I abstain from developing the more accurate formulæ for determining these constants from the sum of the squares of the distances, or from the sum of the distances (the string), also in the case of stray shots, and give in conclusion the most probable shots of the marksmen A and B. They are,

$$\text{For marksman A, } \epsilon_a = \frac{5}{\sqrt{(2l \frac{1.0.0}{6.4})}} = 5.292.$$

$$\text{For marksman B, } \epsilon_b = \frac{10}{3\sqrt{(2l \frac{1.0.0}{3.6})}} = 4.664.$$

PROBLEM 436.—“Integrate the equation

$$x^m y^n (aydx + bxdy) = x^{m'} y^{n'} (a'ydx + b'xdy).”$$

SOLUTION BY PROF. W. W. BEMAN.—Evidently $x^{-m-1}y^{-n-1}$ is an integrating factor of the first member, whence an integral of $x^m y^n (aydx + bxdy) = 0$, is $\log(x^a y^b) = c$, or $x^a y^b = C$.

The general form of integrating factor of the first member is, then,

$$x^{-m-1}y^{-n-1} \phi(x^a y^b).$$

In the same way we may obtain a general integrating factor for the second member,

$$x^{-m'-1}y^{-n'-1} \phi(x^a y^b).$$

That these two may be equal, we must have

$$x^{m'+1}y^{n'+1} \phi(x^a y^b) = x^{m+1}y^{n+1} \phi(x^a y^b).$$

Let $\phi(x^a y^b) = (x^a y^b)^r$, $\phi(x^a y^b) = (x^a y^b)^s$, r and s being indeterminate. Then

$$x^{m'+1}y^{n'+1} (x^a y^b)^r = x^{m+1}y^{n+1} (x^a y^b)^s.$$

$$\therefore m' + ar = m + a's, \quad n' + br = n + b's;$$

$$r = \frac{a'(n-n')-b'(m-m')}{a'b-ab'}, \quad s = \frac{a(n-n')-b(m-m')}{a'b-ab'}.$$

The integrating factor becomes

$$x^{ar-m-1}y^{br-n-1} = x^{a's-m'-1}y^{b's-n'-1};$$

$$\therefore x^{ar-1}y^{br-1}(aydx + bxdy) = x^{a's-1}y^{b's-1}(a'ydx + b'xdy).$$

$$\text{Integrating, } \frac{1}{r} x^{ar} y^{br} = \frac{1}{s} x^{a's} y^{b's} + C.$$

When $a'b = ab'$ the equation is immediately integrable.

This solution is based upon that of a special form of the equation given, found in Hoüel's Calcul Infinitesimal.